

BRCA1/BRCA2: CanVIG-UK Gene-Specific Guidance



Date: 14/02/2025 Version: 1.22

A. Garrett^{1,2}, S. Allen¹, L. Loong¹, M Durkie³, G.J. Burghel⁴, R. Robinson⁵, A. Callaway⁶, J. Field⁷, B. Frugtniet², S. Palmer-Smith⁸, J. Grant⁹, J. Pagan¹⁰, T. McDevitt¹¹, T. McVeigh¹², H. Hanson^{1,13,14}, C. Turnbull^{1,11}

- 1) Division of Genetics and Epidemiology, The Institute of Cancer Research, London, UK.
- 2) St George's University Hospitals NHS Foundation Trust, Tooting, London, UK
- 3) Sheffield Diagnostic Genetics Service, NEY Genomic Laboratory Hub, Sheffield Children's NHS Foundation Trust, Sheffield, UK
- 4) Manchester Centre for Genomic Medicine and NW Laboratory Genetics Hub, Manchester University Hospitals NHS Foundation Trust, Manchester, UK
- 5) Yorkshire Regional Genetics Service, Leeds Teaching Hospitals NHS Trust, Leeds, UK
- 6) Wessex Regional Genetics Laboratory, Salisbury NHS Foundation Trust, Salisbury, UK
- 7) Genomics and Molecular Medicine Service, Nottingham University Hospitals NHS Trust, Nottingham, UK
- 8) Institute of Medical Genetics, University Hospital of Wales, Cardiff and Vale University Health Board, Cardiff, UK
- 9) Laboratory Genetics, Queen Elizabeth University Hospital, NHS Greater Glasgow and Clyde, Glasgow, UK
- 10) South East Scotland Clinical Genetics, Western General Hospital, Edinburgh, UK.
- 11) Department of Clinical Genetics, CHI at Crumlin, Dublin, Ireland
- 12) The Royal Marsden NHS Foundation Trust, Fulham Road, London
- 13) Peninsula Regional Genetics Service, Royal Devon University Healthcare NHS Foundation Trust, Exeter, UK
- 14) Faculty of Health and Life Sciences, University of Exeter, Exeter, UK

For use in conjunction with CanVIG-UK Consensus Specification for Cancer susceptibility Genes of ACGS Best Practice Guidelines for Variant Classification. Evidence lines for which there are no gene-specific recommendations should be reviewed in context of CanVIG-UK Consensus Specification for Cancer Susceptibility Genes.

Evidence towards Pathogenicity

Evidence element and evidence strengths allowed	Thresholds/data-sources/applications specifically relevant to BRCA1/BRCA2
PS4: Case-control: The prevalence of the variant in affected individuals is significantly increased compared with the prevalence in controls	<div style="display: flex; align-items: center;"> <div style="margin-right: 10px;"> <div style="background-color: red; color: white; padding: 2px 5px;">_VSTR</div> <div style="background-color: orange; color: white; padding: 2px 5px;">_STR</div> <div style="background-color: yellow; color: black; padding: 2px 5px;">_MOD</div> <div style="background-color: green; color: white; padding: 2px 5px;">_SUP</div> </div> <div> NHSD case control data can be used for case-control analysis: <ul style="list-style-type: none"> Controls should represent appropriate ethnicity and sex matching (i.e. female individuals from UKBiobank should be used if the case series consists predominantly of females, as with the current NHSD case series) For unenriched cases, an OR threshold of >4 should be used based on the ENIGMA threshold for high-penetrance genes. However, as this is an enriched series, a dataset-specific enrichment factor should be used to calculate the odds ratio (OR) threshold where available. Otherwise, an OR threshold of >8 should be used Current data/denominator counts for base substitutions are available at CanVar-UK For non-base-substitutions i.e. deletions/duplications/insertions, NHSD counts can be accessed from CanVIG-UK <p>If there are insufficient data to perform case-control analyses, PS4 can be applied:</p> <ul style="list-style-type: none"> at PS4_sup if there are observations of the variant in ≥5 different families and the variant is seen in ≤ 1/50,000 individuals in UKBiobank at PS4_mod if there are observations of the variant in ≥10 different families and the variant is absent from UKBiobank. Families used must have a pattern of diagnoses consistent with a hereditary breast and ovarian cancer syndrome. </div> </div>
PM2: Absent from controls (or at extremely	<div style="display: flex; align-items: center;"> <div style="margin-right: 10px;"> <div style="background-color: yellow; color: black; padding: 2px 5px;">_MOD</div> <div style="background-color: green; color: white; padding: 2px 5px;">_SUP</div> </div> <div> Female controls of any/all ethnicities from gnomAD v4.1 (or the non-UKBiobank partition of gnomAD v4.1 if using UKBiobank for </div> </div>

low frequency if recessive) in ESP, 1000GP, or ExAC		PS4) should be used (due to low penetrance in male pathogenic variant carriers). If the variant is absent from non-UKBiobank female controls but is present in UKBiobank female controls, then PM2 may be applied at a maximum of supporting. Otherwise, the main CanVIG-UK consensus guidance should be followed.
PVS1: Predicted null variant (in a gene where LOF is a known mechanism of disease)	_VSTR _STR _MOD _SUP	Please refer to the ENIGMA <i>BRCA1</i> and <i>BRCA2</i> VCEP look-up table (Specifications Table 4: "Summary of codes applicable for variants considered against the <i>BRCA1</i> and <i>BRCA2</i> PVS1 decision trees") for the latest advice on application of PVS1 and PM5_PTC for variants across <i>BRCA1</i> and <i>BRCA2</i> .
PS1: Same amino acid change as an established variant	_STR	Within forthcoming ACMG guidance, it is anticipated that these elements will all be incorporated within PP3 and only awarded to variants within key domains.
PM4: Protein-length-changing variant	_MOD _SUP	In the interim, we recommend:
PP3: In silico: Multiple lines of computational evidence support a deleterious effect on the gene or gene product	_SUP	<ul style="list-style-type: none"> Use of PM1_sup and/or PM4_sup for any variant within <i>BRCA1</i> RING (aa 2-101), BRCT (aa 1650-1857) COILED-COIL DOMAIN (aa 1391-1424), <i>BRCA2</i> DNA-binding domain (aa 2481-3186), and <i>BRCA2</i> PALB2 binding domain (aa 10-40).
PM5: Novel missense change at an amino acid residue where a different missense change determined to be pathogenic seen before	_MOD _SUP	<ul style="list-style-type: none"> Use of PM1_mod or PM4_mod for a variant at specific residues¹: RING: 18, 22, 37, 39, 41, 44, 47, 61, 64, 71 BRCT: 1685, 1688, 1697, 1699, 1706, 1708, 1715, 1736, 1738, 1739, 1748, 1764, 1766, 1770, 1775, 1786, 1837, 1838, 1839, 1853 DBD: 2607, 2626, 2627, 2663, 2722, 2723, 2748, 3052, 3124
PM1, PP2: Enrichment/constraint: PP2: Missense variant in a gene that has a low rate of benign missense variation and in which missense variants are a common mechanism of disease PM1: Located in a mutational hot spot and/or critical and well-established functional domain (e.g. active site of an enzyme) without benign variation	_STR _MOD _SUP	<ul style="list-style-type: none"> PM1 cannot be used where functional data are being used for PS3, as per main CanVIG-UK guidance PP2 should not be used for <i>BRCA1/BRCA2</i> Use PM5_PTC as per the ENIGMA <i>BRCA1</i> and <i>BRCA2</i> VCEP guidelines Use PM5, PS1, PP3 otherwise as per CanVIG-UK Consensus Specification
PS3: Functional: Well-established in vitro or in vivo functional studies supportive of a damaging effect on the gene or gene product	_VSTR _STR _MOD _SUP	<p><u>BRCA1</u> Strong: Findlay et al, 2018²; Bouwman et al, 2020³; Starita et al, 2018⁴ Supporting: Fernandes et al, 2019⁵; Petitalot et al, 2019⁶</p> <p><u>BRCA2:</u> Strong: *Guidugli et al, 2018⁷; *Hart et al, 2019⁸; *Richardson et al, 2021⁹; Ikegami et al, 2020¹⁰; *Hu et al, 2022¹¹; *Hu et al, 2024¹², Biswas et al, 2020¹³ Moderate: Mesman et al, 2019¹⁴ *Please note that results from these studies utilise the same assay, and as such results should not be used in combination to attain higher evidence scores for PS3 or BS3.</p> <p><u>Additional Notes:</u> See the full list of CanVIG-UK reviewed functional assays and scores on the CanVIG-UK website. See the table at the bottom of this document for guidance on combining assay results.</p>
	_VSTR	

<p>PP1: Co-segregation with disease in multiple affected family members in a gene definitively known to cause the disease</p>	<p>_STR _MOD _SUP</p>	<p>Segregation evidence from multifactorial analysis data is incorporated within the combined scores described in the PP4/BP5 recommendations.</p> <p>Meiosis counting approaches may be used in addition if this evidence comes from families not already included in the multifactorial analyses. Evidence cannot exceed ‘Very strong’</p>
<p>PS2/PM6: De novo (maternity and paternity confirmed/unconfirmed) in a patient with the disease and no family history</p>	<p>_STR _MOD _SUP</p>	
<p>PM3: in trans with a pathogenic variant</p>	<p>_STR _MOD _SUP</p>	<p>Frequency data regarding co-occurrence in trans is incorporated within the combined scores described in the PP4/BP5 recommendations.</p> <p>In addition, the SVI recommendations for in trans Criterion (PM3) can be used for either BRCA1 or BRCA2 for individuals with a Fanconi anaemia phenotype if this evidence comes from families not already included in the multifactorial analyses used for PP4. Evidence towards a Fanconi phenotype comprise:</p> <ul style="list-style-type: none"> • Clinical: diagnosis of childhood cancer or skeletal/structural/developmental abnormalities • Molecular/Cellular: aberration on mitomycin-induced chromosomal breakage +/- depletion of BRCA2 in lymphocytes <p>Both clinical and molecular/cellular aberrations must be present for a case to contribute to evidence Evidence cannot exceed ‘Strong’</p> <p>Note: Caution is required in inferring the pathogenicity for the monoallelic phenotype, as variants may be hypomorphic (e.g. a variant contributing and causing a Fanconi anaemia phenotype may be low penetrance for breast cancer). Where the majority of evidence for variant pathogenicity comes from observations of the variant in cases of Fanconi Anaemia, it may be appropriate to comment on this in the clinical report.</p>
<p>PP4: Phenotypic specificity (Patient’s phenotype or family history is highly specific for a disease with a single genetic aetiology)</p>	<p>_VSTR _STR _MOD _SUP</p>	<p>Published multifactorial analysis data providing likelihood ratios (LR) or log likelihood ratios (LLR) encompassing multiple evidence types can be applied under PP4/BP5. The combined score should be used, representing the totality of evidence.</p> <p>Suitable analyses:</p> <ul style="list-style-type: none"> • Easton et al, 2007¹⁵ • Vallée et al, 2012¹⁶ • Parsons et al, 2020¹⁷ • Caputo et al, 2021¹⁸ • Li et al, 2020¹⁹ <p>Evidence is presented as either a Likelihood Ratio (LR) or Log Likelihood Ratio (LLR).</p> <p>If evidence is supplied as an LR: Use the table below to directly convert the LR to the applicable Evidence Strength.</p> <p>If evidence is supplied as an LLR: First, convert the LLR to a Likelihood Ratio (LR) by finding the exponent of the LLR; conversion of an LLR to an LR can be done using the =EXP(LLR) function within Excel if a natural log has been used or the =10^LLR function in Excel if log to base 10 has been used.</p>

Once the LR is calculated, use the table below to directly convert the LR to the applicable Evidence Strength (LR of 7.38 = MOD).

Conversions from LR or LLR to Evidence (Exponent) points is also available for applicable variants at <https://canvaruk.org/>, where 'ACMG LLR' is equivalent to Evidence (Exponent) points.

Likelihood Ratio	Evidence (Exponent) Points	Evidence Strength towards pathogenicity
2.08 – 4.30	1	SUP
4.31 – 18.70	2	MOD
18.71 – 350.40	4	STR
≥ 350.41	8	VSTR

Explanatory Notes:

- Where multiple potentially valid LR/LLRs are available for a variant, the value from the most recent publication should be used.
- Evidence (Exponent) Points are calculated by applying the logarithm of the LR to base 2.08. A calculated Evidence Point that is between two categories (eg 3 points) is assigned the weaker strength of the two categories it lies between (eg MOD for 3 points)

Evidence towards Benignity

BA1/BS1: Allele frequency is “too high” in ExAC or gnomAD for disorder	_SA _STR	<p>BA1: MTAF = 0.001 (0.1%) BS1: MTAF = 0.0001 (0.01%)</p> <p>The MTAF (maximum tolerated allele frequency) has been calculated using cardiodb using the calculate AF function: prevalence 0.125; genetic heterogeneity 0.01; allelic heterogeneity 1 (BA1) 0.1 (BS1); penetrance 0.72 (BRCA1), 0.69 (BRCA2). See training resources from Miranda Durkie for further details.</p> <p>Female controls should be used when determining the maximum allele count / filtering allele frequency.</p> <p>See consensus guidelines for further details on Grpmax Filtering AF, and the use of cardiodb for calculating the maximum allele count / filtering allele frequency.</p>
BS2: Observation in controls inconsistent with disease penetrance. Observed in a healthy adult individual for a recessive (homozygous), dominant (heterozygous), or X-linked (hemizygous) disorder, with full penetrance expected at an early age	_STR _SUP	
BP4: In silico: Multiple lines of computational evidence suggest no impact on gene or gene product (conservation, evolutionary, splicing impact, etc.)	_SUP	
BP1: Missense variant in a gene for which primarily truncating variants are known to cause disease	_SUP	Can be used for missense variants with no predicted splicing effect (as per main CanVIG-UK consensus specification) at non-conserved residues outside of <i>BRCA1</i> RING (aa 2-101), BRCT (aa 1650-1857) COILED-COIL DOMAIN (aa 1391-

		1424) and <i>BRCA2</i> DNA-binding domain (aa 2481-3186) and <i>BRCA2 PALB2</i> binding domain (aa 10-40)															
BP7: Synonymous (silent) variant for which splicing prediction algorithms predict no impact to the splice consensus sequence	_SUP																
BP3: In-frame deletions/insertions in a repetitive region	_SUP																
BS3: Well-established <i>in vitro</i> or <i>in vivo</i> functional studies show no damaging effect on protein function or splicing	_STR _MOD _SUP	*see PS3															
BS4: Non segregation with disease	_STR _SUP	*see PP1															
BP2: Observed in trans with a pathogenic variant for a fully penetrant dominant gene/disorder or observed in cis	_STR _SUP	*see PM3															
BP5: Alternate molecular basis for disease	_VSTR _STR _MOD _SUP	*see PP4 for explanation															
		<table border="1"> <thead> <tr> <th>Likelihood Ratio</th> <th>Evidence (Exponent) Points</th> <th>Evidence Strength towards benignity</th> </tr> </thead> <tbody> <tr> <td>0.48 – 0.23</td> <td>-1</td> <td>SUP</td> </tr> <tr> <td>0.22 – 0.05</td> <td>-2</td> <td>MOD</td> </tr> <tr> <td>0.049 – 0.00285</td> <td>-4</td> <td>STR</td> </tr> <tr> <td><0.00284</td> <td>-8</td> <td>VSTR</td> </tr> </tbody> </table>	Likelihood Ratio	Evidence (Exponent) Points	Evidence Strength towards benignity	0.48 – 0.23	-1	SUP	0.22 – 0.05	-2	MOD	0.049 – 0.00285	-4	STR	<0.00284	-8	VSTR
Likelihood Ratio	Evidence (Exponent) Points	Evidence Strength towards benignity															
0.48 – 0.23	-1	SUP															
0.22 – 0.05	-2	MOD															
0.049 – 0.00285	-4	STR															
<0.00284	-8	VSTR															

Recommendations for the management of conflicting functional assay results See table below for management of discrepancy for *BRCA1* variants between Findlay et al, 2018² and Bouwman et al, 2020³ discordant assay results. For more general guidance regarding conflicting results from other functional assays, refer to the table in the main CanVIG-UK consensus specification.

Findlay Class	Findlay Score	Bouwman Platinum	Bouwman Olaparib	Bouwman DR-GFP	PS3_STR	BS3_STR
LOF	<-1.328	All deleterious/ likely deleterious (1 intermediate allowed)			✓	✗
LOF	<-1.328	Any are neutral/likely neutral			✗	✗
INT (towards LOF)	-1.328 to -1.038	All deleterious/ likely deleterious			✓	✗
INT (towards FUNC)	-1.038 to -0.748	All neutral/likely neutral			✗	✓
INT	-1.328 to -0.78	Conflicting results or any intermediate			✗	✗
FUNC	>-0.748	All neutral/likely neutral (1 intermediate allowed)			✗	✓
FUNC	>-0.748	Any are deleterious/likely deleterious			✗	✗

N.B: Bouwman et al, 2020 “not clear” refers to opposite categorisation \pm the standard deviation of repeat experiments and should be treated as conflicting assay results. Where a variant is LOF on the Findlay et al assay and has an RNA score of <-2, this indicates that LOF is due to interference with splicing and therefore should not be treated as conflicting evidence if the variant is neutral on the Bouwman et al assay.

Version History/Amendments

Revised version	Date	Section	Update	Amended by	Approved by
1.12	01/09/2021	PP4	Guidance on use of LLRs from published epidemiological studies amended to account for the use of natural logs in the statistics presented	Garrett	Turnbull
1.12	01/09/2021	PM1	Addition of critical residues in the DNA binding domain of BRCA2. Critical residues in all listed functional domains updated to mirror draft 2021 ENIGMA guidance	Garrett	Turnbull
1.12	01/09/2021	BP1	Resolution of typo in BRCT region specification	Garrett	Turnbull
1.13	15/10/2021	PVS1	Clarification that PVS1 may not be applicable for some of the variants at ENIGMA specified positions	Garrett	Turnbull
1.14	02/12/2021	PS4	Terminology change to reflect transition of PHE to NHSD	Garrett	Turnbull
1.14	02/12/2021	PVS1	Addition of recommendations for variants within the first 100bp	Callaway	CStAG
1.15	28/04/2022	PM1/ PM4	Clarification that PM1 _{sup} and PM4 _{sup} may be used in combination but PM4 not to be used at moderate. Removal of mention that CanVIG-UK provide non-white ethnicity counts under PS4	Garrett	CStAG
1.16	28/07/2022	PP5	Removal of Lindor <i>et al</i> 2011 paper from recommended genetic epidemiology papers to use in calculating Evidence (Exponent) Points.	Allen	Turnbull
1.17	27/09/2022	PS4	Addition of PS4 _{mod} application where ≥10 HBOC families observed.	Garrett	CStAG
1.18	31/03/2023	PP5/ BP6	Rewording of application details for clarity. Removal of reputable source evidence from PP5 per consensus specification. Addition of recommended analyses papers.	Allen	CStAG
1.18	31/03/2023	PVS1/ PM1/ PS3	Incorporation of functional assays, hotspot, and splice sites from upcoming ENIGMA recommendations.	Allen	CStAG
1.19	26/05/2023	PS4/PM2/ BA1/BS1	Update of databases to be used in-line with consensus specification.	Garrett	CStAG
1.19	27/05/2023	PS4	Update on case-counting approach where variant seen in multiple cases but also observed in control datasets.	Garrett	CStAG
1.19	15/09/2023	BA1/BS1	Clarification of MTAF usage and use of the filtering allele frequency.	Callaway	CStAG
1.19	28/09/2023	PP4/BP5	Moved multifactorial evidence from PP5/BP6 to PP4/BP5 in alignment with ENIGMA. PP5/BP6 evidence code removed.	CStAG	CStAG
1.19	29/09/2023	PM4	Added application at 'Moderate' for in-frame in/dels at specific residues.	Allen	CStAG
1.19	29/09/2023	PP1/PM3/ PP4/BP5	Recommendation to use the combined multifactorial score under PP4/BP5, rather than individual subcomponent scores	Garrett	CStAG
1.19	29/09/2023	PS4	Change of OR threshold from >10 to >8 for enriched case series where dataset-specific enrichment factors are not	CStAG	CStAG

			available (eg variant count releases from NHSD/NDRS)		
1.19	29/09/2023	PVS1	Update of NMD boundary as per ENIGMA VCEP BRCA1 and BRCA2 guidelines	CStAG	CStAG
1.20	24/01/2024	PS3/BS3	Added Ikegami et al 2020 and Hu et al 2022 papers to functional study review list (PS3/BS3)	Allen	CStAG
1.20	24/01/2024	PVS1	Update to refer use of PVS1 to ENIGMA VCEP	Allen	CStAG
1.20	30/04/2024	PM2	Replaced ref to cancer-free gnomAD v2.1.1 and UKBiobank with gnomAD v4.1, clarified application of PM2 strength where data is in UKBiobank but absent from other gnomAD datasets.	Allen	CStAG
1.20	30/04/2024	PS3/BS3	Added statement to highlight assay result overlap for Couch lab assays	Allen	CStAG
1.20	30/04/2024	PS3/BS3	Updated functional assays scoring link	Allen	CStAG
1.20	07/05/2024	PM3	Typing error amendment – ‘PP5’ to ‘PP4’	Allen	CStAG
1.21	25/07/2024	PVS1	Removed splice tables to refer only to the ENIGMA VCEP guidelines	Allen	CStAG
1.22	28/01/2025	PP4/BP5	Added Li <i>et al.</i> , 2020 as reference for suitable studies	Allen	Turnbull

References

1. ENIGMA. *BRCA1/2 Gene Variant Classification Criteria Version 2.5.1 2017* [Available from: <https://enigmaconsortium.org/library/general-documents/enigma-classification-criteria/>].
2. Findlay GM, Daza RM, Martin B, et al. Accurate classification of BRCA1 variants with saturation genome editing <https://sge.gs.washington.edu/BRCA1/>. *Nature* 2018;562(7726):217-22. doi: 10.1038/s41586-018-0461-z [published Online First: 2018/09/14]
3. Bouwman P, van der Heijden I, van der Gulden H, et al. Functional Categorization of BRCA1 Variants of Uncertain Clinical Significance in Homologous Recombination Repair Complementation Assays. *Clinical cancer research : an official journal of the American Association for Cancer Research* 2020;26(17):4559-68. doi: 10.1158/1078-0432.Ccr-20-0255 [published Online First: 2020/06/18]
4. Starita LM, Islam MM, Banerjee T, et al. A Multiplex Homology-Directed DNA Repair Assay Reveals the Impact of More Than 1,000 BRCA1 Missense Substitution Variants on Protein Function. *Am J Hum Genet.* 2018;103(4):498-508. doi:10.1016/j.ajhg.2018.07.016 [published Online First: 2018/09/12]
5. Fernandes VC, Golubeva VA, Di Pietro G, et al. Impact of amino acid substitutions at secondary structures in the BRCT domains of the tumor suppressor BRCA1: Implications for clinical annotation. *The Journal of biological chemistry* 2019;294(15):5980-92. doi: 10.1074/jbc.RA118.005274 [published Online First: 2019/02/16]
6. Petitalot A, Dardillac E, Jacquet E, et al. Combining Homologous Recombination and Phosphopeptide-binding Data to Predict the Impact of BRCA1 BRCT Variants on Cancer Risk. *Mol Cancer Res* 2019;17(1):54-69. doi: 10.1158/1541-7786.Mcr-17-0357 [published Online First: 2018/09/28]
7. Guidugli L, Shimelis H, Masica DL, et al. Assessment of the Clinical Relevance of BRCA2 Missense Variants by Functional and Computational Approaches. *American journal of human genetics* 2018;102(2):233-48. doi: 10.1016/j.ajhg.2017.12.013 [published Online First: 2018/02/06]
8. Hart SN, Hoskin T, Shimelis H, et al. Comprehensive annotation of BRCA1 and BRCA2 missense variants by functionally validated sequence-based computational prediction models. *Genetics in medicine : official journal of the American College of Medical Genetics* 2019;21(1):71-80. doi: 10.1038/s41436-018-0018-4 [published Online First: 2018/06/10]
9. Richardson ME, Hu C, Lee KY, et al. Strong functional data for pathogenicity or neutrality classify BRCA2 DNA-binding-domain variants of uncertain significance. *American journal of human genetics* 2021;108(3):458-68. doi: 10.1016/j.ajhg.2021.02.005 [published Online First: 2021/02/21]

10. Ikegami M, Kohsaka S, Ueno T, et al. High-throughput functional evaluation of BRCA2 variants of unknown significance. *Nat Commun.* 2020;11(1):2573. doi:10.1038/s41467-020-16141-8 [published Online First: 2020/05/22]
11. Hu C, Susswein LR, Roberts ME, et al. Classification of BRCA2 Variants of Uncertain Significance (VUS) Using an ACMG/AMP Model Incorporating a Homology-Directed Repair (HDR) Functional Assay. *Clin Cancer Res.* 2022;28(17):3742-3751. doi:10.1158/1078-0432.CCR-22-0203 [published Online First: 2022/09/01]
12. Hu C, Huang H, Na J, et al. Functional analysis and clinical classification of 462 germline BRCA2 missense variants affecting the DNA binding domain. *Am J Hum Genet.* 2024;111(3):584-593. doi:10.1016/j.ajhg.2024.02.002 [published Online First: 2024/02/27]
13. Biswas K, Lipton GB, Stauffer S, et al. A computational model for classification of BRCA2 variants using mouse embryonic stem cell-based functional assays. *NPJ Genom Med.* 2020;5(1):52. Published 2020 Dec 8. doi:10.1038/s41525-020-00158-5
14. Mesman RLS, Calléja FMGR, Hendriks G, et al. The functional impact of variants of uncertain significance in BRCA2. *Genet Med.* 2019;21(2):293-302. doi:10.1038/s41436-018-0052-2 [published Online First: 2018/07/10]
15. Easton DF, Deffenbaugh AM, Pruss D, et al. A systematic genetic assessment of 1,433 sequence variants of unknown clinical significance in the BRCA1 and BRCA2 breast cancer-predisposition genes. *American journal of human genetics* 2007;81(5):873-83. doi: 10.1086/521032 [published Online First: 2007/10/10]
16. Vallée MP, Francy TC, Judkins MK, et al. Classification of missense substitutions in the BRCA genes: a database dedicated to Ex-UVs. *Hum Mutat.* 2012;33(1):22-28. doi:10.1002/humu.21629 [published Online First: 2011/11/03]
17. Parsons MT, Tudini E, Li H, et al. Large scale multifactorial likelihood quantitative analysis of BRCA1 and BRCA2 variants: An ENIGMA resource to support clinical variant classification. *Human mutation* 2019;40(9):1557-78. doi: 10.1002/humu.23818 [published Online First: 2019/05/28]
18. Caputo SM, Golmard L, Léone M, et al. Classification of 101 BRCA1 and BRCA2 variants of uncertain significance by cosegregation study: A powerful approach. *Am J Hum Genet.* 2021;108(10):1907-1923. doi:10.1016/j.ajhg.2021.09.003 [published Online First: 2021/09/30]
19. Li H, LaDuca H, Pesaran T, et al. Classification of variants of uncertain significance in BRCA1 and BRCA2 using personal and family history of cancer from individuals in a large hereditary cancer multigene panel testing cohort. *Genet Med.* 2020;22(4):701-708. doi:10.1038/s41436-019-0729-1